

Figure 1. i.v injection of Vitamin C provides optimal, TET2-dependent anti-PD-L1 immunotherapy efficacy.



TET2-KO clones of CT-26 cells made by CRISPR/Cas9 system were confirmed by western blotting

Figure 1C


Figure 1. i.v injection of Vitamin C provides optimal, TET2-dependent anti-PD-L1 immunotherapy efficacy.



TET2-KO clones of CT-26 cells made by CRISPR/Cas9 system were confirmed by western blotting

Figure 1C


Figure 6. IFN-γ signal communication was identified in the WT tumor microenvironment and played a critical role in regulating MHC I gene transcription.



Protein Immunoprecipitation (IP) was conducted between TET2 and STAT1 in the presence of IFNg or VC treatment.

Figure 6D


Figure 6. IFN-γ signal communication was identified in the WT tumor microenvironment and played a critical role in regulating MHC I gene transcription.



Protein Immunoprecipitation (IP) was conducted between TET2 and STAT1 in the presence of IFNg or VC treatment.

Figure 6D


Figure 6. IFN-γ signal communication was identified in the WT tumor microenvironment and played a critical role in regulating MHC I gene transcription.



Protein Immunoprecipitation (IP) was conducted between TET2 and STAT1 in the presence of IFNg or VC treatment.

Figure 6D


Figure 6. IFN-γ signal communication was identified in the WT tumor microenvironment and played a critical role in regulating MHC I gene transcription.



Protein Immunoprecipitation (IP) was conducted between TET2 and STAT1 in the presence of IFNg or VC treatment.

Figure 6D


Figure 6. IFN-γ signal communication was identified in the WT tumor microenvironment and played a critical role in regulating MHC I gene transcription.



Protein Immunoprecipitation (IP) was conducted between TET2 and STAT1 in the presence of IFNg or VC treatment.

Figure 6D


Figure 6. IFN-γ signal communication was identified in the WT tumor microenvironment and played a critical role in regulating MHC I gene transcription.



Protein expression of MHC I antigen-presenting genes TAP1, TAPBP was determined by Western-blot in the WT or TET2-KO B16-OVA cells under IFNg treatment or STAT1 siRNA knock down.

Figure 6F


Figure 6. IFN-γ signal communication was identified in the WT tumor microenvironment and played a critical role in regulating MHC I gene transcription.



Protein expression of MHC I antigen-presenting genes TAP1, TAPBP was determined by Western-blot in the WT or TET2-KO B16-OVA cells under IFNg treatment or STAT1 siRNA knock down.


Figure 6F

Figure 6. IFN-γ signal communication was identified in the WT tumor microenvironment and played a critical role in regulating MHC I gene transcription.



Protein expression of MHC I antigen-presenting genes TAP1, TAPBP was determined by Western-blot in the WT or TET2-KO B16-OVA cells under IFNg treatment or STAT1 siRNA knock down.

Figure 6. IFN-γ signal communication was identified in the WT tumor microenvironment and played a critical role in regulating MHC I gene transcription.



Protein expression of MHC I antigen-presenting genes TAP1, TAPBP was determined by Western-blot in the WT or TET2-KO B16-OVA cells under IFNg treatment or STAT1 siRNA knock down.

Figure 6F

Figure 6. IFN-γ signal communication was identified in the WT tumor microenvironment and played a critical role in regulating MHC I gene transcription.



Protein expression of MHC I antigen-presenting genes TAP1, TAPBP was determined by Western-blot in the WT or TET2-KO B16-OVA cells under IFNg treatment or STAT1 siRNA knock down.

Figure 6F